

California Bioscience

Product Datasheet

Product Name	Glomerular Basal Membrane Human Recombinant
Cata No	CB501008
Source	Sf9 insect cells.
Synonyms	Collagen alpha-3(IV) chain, Goodpasture antigen, COL4A3, Glomerular Basal Membrane, GBM.

Description

Type IV collagen is a major structural component of basement membranes. It is a multimeric protein composed of 3 alpha subunits, which are encoded by 6 different genes, alpha 1 through alpha 6. Each of these alpha subunits can form a triple helix structure with 2 other subunits to form type IV collagen. The Goodpasture syndrome is a condition in which autoantibodies bind to the collagen molecules in the basement membranes of alveoli and glomeruli. The epitopes that elicit these autoantibodies are restricted basically to the non-collagenous C-terminal domain of the protein. There are numerous alternative transcripts that appear to be unique to the human COL4A3 and alternative splicing is limited to the six exons that encode this C-terminal domain. COL4A3 is also linked to an autosomal recessive form of Alport syndrome. The mutations contributing to the Alport syndrome are also situated within the exons that encode this C-terminal region. COL4A3 is organized in a head-to-head conformation with another type IV collagen gene so that each gene pair shares a common promoter. Several exons of COL4A3 are interspersed with exons of an uncharacterized gene which is on the opposite strand.

Human α3 chain of collagen IV; identical with the antigen called "glomerular basal membrane antigen" (GBM). Recombinant antigen for solid (ELISA) and fluid phase diagnostic assays. Binds IgG-type human auto-antibodies. Calculated Molecular weight: 43,591 Dalton. Calculated isoelectric point: pH 8.9.

cDNA coding for a minicollagen version of the human collagen IV α3 chain fused to a hexa-histidine purification tag. The term minicollagen designates the removal of most of the epitope-less triplehelical collagenous region (situated between the N-terminal 7S domain and the C-terminal noncollagenous NC1 domain), which is a requirement for recombinant production of this antigen.

Physical Appearance

Sterile Filtered clear solution.

Formulation

The protein solution (0.3 mg/ml) contains 20mM Hepes, pH 8.0, 80mM NaCl, 20% Glycerol.

Stability

Recommendations for storage buffer: ion strength 100mM, neutral to slightly alkaline pH and 20 % glycerol as cryoprotective agent. Storage temperature: -70°to -80°C. **Please prevent freeze-thaw cycles.**

Applications

Standard ELISA test Immuno-Dot test with positive/negative sera panels. SDS-PAGE Western-Blot



Product Datasheet

* For Non-Clinical Research Use Only *